

D-125 Efficacy Overview

EPA Registration # 61178-1
Revised Date: January 27, 2009

1. *Acinetobacter calcoaceticus var anitratus*: causes bacterial meningitis, fulminating septicaemia, pulmonary and ophthalmic infections, chronic synovitis (joint pain/inflammation), skin diseases, wound infections and postoperative urinary tract infections.

Category and/or Source: Gram negative clinical isolate **Contact Time:** 10 minutes

2. *Acinetobacter calcoaceticus var lwoffii*: causes bacteremia, pneumonia, meningitis, abdominal inflammation, endocarditis, and infections of the urinary tract and skin.

Category and/or Source: Gram negative clinical isolate **Contact Time:** 10 minutes

3. *Actinobacillus pleuropneumoniae*: causes a bacterial upper respiratory disease in pigs, resulting in lethargy, cough, and other breathing difficulties. The organism is most problematic in intensive pig production operations.

Category and/or Source: ATCC 27088 **Contact Time:** 10 minutes

4. *Actinomyces pyogenes*: causes severe bacterial mastitis in cattle, characterized by thick, purulent (pus) secretion.

Category and/or Source: ATCC 19411 **Contact Time:** 10 minutes

5. *Adenovirus type 2*: causes nonspecific viral respiratory illness, diarrhea, conjunctivitis (eye inflammation), cystitis, and rashes.

Category and/or Source: ATCC VR-846 **Contact Time:** 10 minutes

6. *Aspergillus candidus*: causes a rare infection of the lungs and is associated with asthma. This fungus is prevalent in the environment.

Category and/or Source: Environmental fungus **Contact Time:** 10 minutes

7-8. *Aspergillus niger*: causes a rare infection of the lungs and is associated with asthma. This fungus is common in the environment. The fungus is characterized by dense growth of black spores.

Category and/or Source: Environmental fungus | AIDS patient isolate **Contact Time:** 10 minutes

9. *Avian Influenza/Turkey Wisconsin Virus*: causes influenza infection of birds. The virus is very similar to the avian influenza H5N1 virus, which is thought to have potential for human crossover as a pandemic strain.

Category and/or Source: ATCC VR-798 **Contact Time:** 10 minutes

10. *Bacillus cereus*: causes gastrointestinal infection and intoxication. The spores of *B. cereus* bacteria commonly contaminate raw foods and food materials, particularly foods that have been in contact with soil. The spores survive cooking and can subsequently germinate and grow under favorable conditions. Consumption of foods contaminated with *B. cereus* may result in disease either by the consumption of pre-formed toxin or by toxins produced by these bacteria in during growth the gut.

Category and/or Source: ATCC 11778 **Contact Time:** 10 minutes


11. *Bacteroides fragilis*: causes various abscesses, mostly in the human gut. These opportunistic anaerobic bacteria may also cause bed sores, pressure sores, aspiration pneumonia, chronic otitis media (ear infection), chronic sinusitis, and osteomyelitis (bone infection).

33 CLINTON ROAD • SUITE 102 • CLINTON SQUARE EXECUTIVE CENTER • WEST CALDWELL, NJ 07006 USA



MICROGEN

Vaccine for the Surface



Category and/or Source: ATCC 43859 **Contact Time:** 10 minutes

12-13. Bordetella bronchiseptica: causes bronchitis in humans and can cause kennel cough in dogs. This bacterium infects the airway, and is closely related to the causative agent of whooping cough, Bordetella pertussis.

Category and/or Source: Gram negative clinical isolate | ATCC 19395 **Contact Time:** 10 minutes

14. Bovine viral diarrhoea virus (BVDV): causes diarrhea in cattle and can cause reproductive problems in pigs. This virus is in the same group of pest viruses as the virus of swine fever (hog cholera).

Category and/or Source: X-800 strain **Contact Time:** 10 minutes

15. Brevibacterium ammoniagenes: causes diaper rash. These bacteria are now known as Corynebacterium ammoniagenes, and are thought to be associated with diaper rash due to their ability to convert urine to ammonia.

Category and/or Source: GBL strain **Contact Time:** 10 minutes

16. Brevundimonas diminuta: causes opportunistic infections and fever. This bacterium is of relatively low clinical significance, but is used frequently to test water filters due to its very small size.

Category and/or Source: Gram negative clinical isolate **Contact Time:** 10 minutes

17. Burkholderia cepacia: causes severe respiratory infections in the immunocompromised. These bacteria also have natural resistance to many antibiotics.

Category and/or Source: Gram negative clinical isolate **Contact Time:** 10 minutes

18. Burkholderia pickettii: causes opportunistic infections in the hospital environment. Also known as Ralstonia pickettii, these bacteria have been isolated from contaminated disinfectant formulations, and are known to infect the blood and tissue around indwelling medical devices. The organism is particularly problematic in patients with cystic fibrosis.

Category and/or Source: ATCC 49729 **Contact Time:** 10 minutes

19. Campylobacter jejuni: causes severe diarrhea. These bacteria cause abdominal pain, nausea, vomiting, diarrhea, and fever. They are found in undercooked meat (especially poultry), unpasteurised milk, and untreated water as a result of contamination by wild fowl. It has been linked with subsequent development of **Guillain-Barré syndrome** (GBS), which usually develops two to three weeks after the initial illness.

Category and/or Source: ATCC 29428 **Contact Time:** 10 minutes

20. Candida albicans: causes opportunistic oral and genital infections in humans. This fungus exists primarily as yeast in the oral cavity, but can infect tissues through the production of invasive filaments called hyphae.

Category and/or Source: AIDS patient isolate **Contact Time:** 10 minutes

21. Canine Coronavirus: causes upper respiratory and gastrointestinal infections in dogs. The virus is related to the human SARS virus.

Category and/or Source: ATCC VR-809 | Strain 1-71 **Contact Time:** 10 minutes

22. Canine Distemper Virus: causes distemper in dogs. This virus is particularly problematic in non-vaccinated populations, including free-living African wild dogs, as well as other carnivores, both free-living and captive.

Category and/or Source: Onderstepoort strain **Contact Time:** 10 minutes


23. Canine Herpesvirus: causes an infection in dogs that can result in various symptoms. The virus is known to cause weakness, depression, discharge from the nose, soft, yellow feces, and a loss of certain motor functions (reflexes). The virus

33 CLINTON ROAD • SUITE 102 • CLINTON SQUARE EXECUTIVE CENTER • WEST CALDWELL, NJ 07006 USA



MICROGEN

Vaccine for the Surface



can also cause keratitis, uveitis, optic neuritis, retinitis, and retinal dysplasia. There is a high mortality rate, approaching 80 percent in puppies less than one week old, and death usually occurs in one to two days.

Category and/or Source: ATCC VR-522 **Contact Time:** 10 minutes

24. Chryseomonas luteola: causes rare opportunistic infections in humans. These bacteria have a propensity to infect hospital patients with health or indwelling medical devices. Most reported cases involve septicemia (blood infection), meningitis (inflammation of nerves or brain tissue), heart infection, or inflammation of the abdominal wall.

Category and/or Source: ATCC 43273 **Contact Time:** 10 minutes

25. Corynebacterium ammoniagenes: causes diaper rash. These bacteria are thought to be associated with diaper rash due to their ability to convert urine to ammonia.

Category and/or Source: ATCC 6872 **Contact Time:** 10 minutes

26. Corynebacterium pseudotuberculosis: causes a severe infection of the lower limbs in horses and cattle. This bacterium is also associated with large, ulcerative skin lesions in about 2-5% of cases.

Category and/or Source: ATCC 19410 **Contact Time:** 10 minutes

27. Cryptococcus neoformans: causes meningitis in the immunocompromised. This fungus recently has also been recognized as a source of pulmonary and general disseminated disease. Many infections with Cryptococcus neoformans are asymptomatic.

Category and/or Source: AIDS patient isolate **Contact Time:** 10 minutes

28. Cytomegalovirus: causes infection of the eyes, throat, and salivary glands. This virus is particularly risky to the immunocompromised, where it can cause a latent infection that further depresses the immune system.

Category and/or Source: ATCC VR-284 **Contact Time:** 10 minutes

29. Enterobacter aerogenes: causes opportunistic, frequently healthcare-associated infections of the skin and skin tissue. The bacterium is problematic because it may become resistant to medical treatments in patients over time.

Category and/or Source: ATCC 13048 **Contact Time:** 1 minute

30-31. Enterobacter agglomerans: causes relatively rare gastrointestinal infections in humans. The bacterium is now called Pantoea agglomerans, and is a recognized plant pathogen.

Category and/or Source: Gram negative clinical isolate | Antibiotic resistant gram negative rod **Contact Time:** 10 minutes

32. Enterobacter cloacae: causes bacteremia, lower respiratory tract infections, skin and soft tissue infections, urinary tract infections, endocarditis (heart infections), intra-abdominal infections septic arthritis, bone infection, and eye infections. This bacterium is most commonly found in healthcare settings, where it is highly associated with invasive medical devices such as catheters.

Category and/or Source: Gram negative clinical isolate **Contact Time:** 10 minutes

33. Enterobacter gergoviae: causes infections associated with indwelling medical devices. These bacteria are rare among Enterobacter infections.

Category and/or Source: Gram negative clinical isolate **Contact Time:** 10 minutes

34. Enterobacter liquefaciens: causes infections associated with indwelling medical devices. These bacteria are rare among Enterobacter infections.


33 CLINTON ROAD • SUITE 102 • CLINTON SQUARE EXECUTIVE CENTER • WEST CALDWELL, NJ 07006 USA

VOICE: 973-575-9025 **FAX:** 973-575-9075 **E-MAIL:** INFO@MICROGENINC.COM
WEB: WWW.MICROGENINC.COM



MICROGEN

Vaccine for the Surface



Category and/or Source: Gram negative clinical isolate **Contact Time:** 10 minutes

35. Enterococcus aerogenes: causes opportunistic infections that are generally associated with the immunocompromised or with indwelling medical devices. These bacteria are of fairly small clinical importance, but are very similar to Enterococcus faecalis, which is of major clinical importance.

Category and/or Source: GBL strain **Contact Time:** 10 minutes

36-37. Enterococcus faecalis: causes opportunistic, but often severe infections of the skin, skin tissues, gastrointestinal tract, and bloodstream. These bacteria are especially problematic in recent years due to their demonstrated propensity to acquire resistance to multiple antibiotics.

Category and/or Source: ATCC 1786-2 VANCOMYCIN resistant -VRE Antibiotic resistant gram positive rod | Gram positive clinical isolate **Contact Time:** 10 minutes

38. Enterococcus faecium: causes opportunistic, but often severe infections of the skin, skin tissues, and bloodstream. These bacteria are very similar to Enterococcus faecalis, and the genus is thought to account for greater than 10% of hospital-acquired infections.

Category and/or Source: ATCC 6569 **Contact Time:** 10 minutes

39. Enterococcus hirae: causes a rare infection of heart valves in humans, as well as other opportunistic infections. It is not as problematic as other members of the genus Enterococcus. These bacteria are not known to become resistant to antibiotics at the current time.

Category and/or Source: ATCC 10541 **Contact Time:** 10 minutes

40. Equine Herpesvirus: causes a respiratory disease of young horses. This virus is primarily associated with coughing, and is thought to require close contact from animal-to-animal for transmission.

Category and/or Source: ATCC VR-700 **Contact Time:** 10 minutes

41. Equine Influenza Virus A: causes a major respiratory disease of horses. Infection with this virus produces flu-like symptoms in horses, but may also prevent horses from drinking for days.

Category and/or Source: ATCC VR-297 **Contact Time:** 10 minutes

42. Escherichia vulneris: causes infection of human wounds. This bacterium was discovered fairly recently, in the early 1980's. Since then, it has also been associated with osteomyelitis (bone infection) and meningitis.

Category and/or Source: Wildtype isolate **Contact Time:** 10 minutes

43 - 46. Escherichia coli: causes a variety of gastrointestinal infections. There are many types of E. coli bacteria, the majority of which are non-pathogenic and live commensally in the gut. The most problematic E. coli are those that produce enterotoxins when growing in the human gut. These extraordinarily powerful toxins act directly on intestinal cells, reversing the flow on ions and causing severe diarrhea. E. coli is also a major hospital pathogen, responsible for greater than 10% of all hospital infections. One particular strain, E. coli O157:H7, causes hemorrhagic intestinal infection and sometimes causes kidney failure.

Category and/or Source: GBL 101 strains | Antibiotic resistant gram negative rod | Gram negative clinical isolate | **Contact Time:** 10 minutes

47. Escherichia coli 0157:H7: causes a severe, hemorrhagic intestinal infection with profuse, bloody diarrhea. These bacteria are commonly found in contaminated ground beef. Once infection is established, they invade intestinal cells and produce toxins that can result in kidney injury. Kidney disease associated with E. coli infection is called Hemolytic Uremic Syndrome (HUS).


Category and/or Source: ATCC 35150 **Contact Time:** 10 minutes

33 CLINTON ROAD • SUITE 102 • CLINTON SQUARE EXECUTIVE CENTER • WEST CALDWELL, NJ 07006 USA



MICROGEN

Vaccine for the Surface



48. Feline Calcivirus: causes a flu-like infection of cats, but is primarily significant because of its similarity to human noroviruses. This virus is recognized by the United States Environmental Protection Agency (USEPA) as a surrogate for noroviruses. Thus, disinfection of feline calcivirus virtually ensures disinfection of human norovirus, which cannot currently be grown or tested in the laboratory.

Category and/or Source: Upjohn Company strain **Contact Time:** 10 minutes

49. Feline Infectious Peritonitis Virus: causes a mild, self-limiting diarrhea in cats. This virus predominantly infects cats that are very young or very old. The virus is thought to be highly transmissible from cats to kittens.

Category and/or Source: ATCC VR-990 **Contact Time:** 10 minutes

50. Flavobacterium meningosepticum: causes meningitis in humans. This bacterium is particularly problematic in children, where infections can be very serious and may result in death.

Category and/or Source: ATCC 10211 **Contact Time:** 10 minutes

51. Haemophilus influenzae: causes bacteremia, and acute bacterial meningitis. It is known as an opportunistic bacterial pathogen. Occasionally, it causes cellulitis, osteomyelitis (bone infection), sore throat, and joint infections. A vaccine (HiB) is available that can prevent infections with this bacteria.

Category and/or Source: ATCC 10211 **Contact Time:** 10 minutes

52. Hafnia alvei: causes diarrhea in humans. This member of the group of bacteria called Enterobacteriaceae is not well understood at this time but is rarely considered to be pathogenic.

Category and/or Source: Gram negative clinical isolate **Contact Time:** 10 minutes

53. HCV (Hepatitis C Virus): causes a blood and fluid-borne infection of the liver in humans. This virus is especially problematic for intravenous illicit drug users and recipients of transfused blood and blood products. No vaccine currently exists for hepatitis C virus, but many are under development. Infection with this virus is associated with decreased liver function and increased likelihood of liver cancer.

Category and/or Source: BVDV Surrogate **Contact Time:** 10 minutes

54. Herpes Simplex Virus type 1: causes small, painful ulcers on the human lips, mouth, and occasionally the ears and genital areas. This virus is known to integrate its DNA into that of the human body and infections are known to occur regularly as cycles. At this time it is not well understood what initiates acute infection or remission. The virus is transmitted by close contact, such as kissing and touching.

Category and/or Source: ATCC VR-260 **Contact Time:** 30 seconds

55. Herpes Simplex Virus type 2: causes small, painful ulcers primarily around the human genital area. This virus is very similar to herpes simplex virus type 1, and symptoms from one virus may often be mistaken for symptoms of the other. The virus is sexually transmitted.

Category and/or Source: ATCC VR-734 **Contact Time:** 30 seconds

56. Human Coronavirus @ 98% Organic Soil Load Tolerance/400 ppm Hard Water: causes gastrointestinal infections in humans and is responsible for about 30% of common colds. This virus is very similar in terms of size and shape to the virus that causes SARS. All age groups can be infected, and severity of infection varies from mild to severe.

Category and/or Source: ATCC VR-740, Strain 229E **Contact Time:** 10 minutes


57. Human Hepatitis B Virus (HBV): causes cirrhosis or liver cancer in humans. The virus is transmitted by contaminated bodily fluids, with the exception of urine, saliva, and stool. Infection of the liver with the virus lasts from one month to many

33 CLINTON ROAD • SUITE 102 • CLINTON SQUARE EXECUTIVE CENTER • WEST CALDWELL, NJ 07006 USA



MICROGEN

Vaccine for the Surface



decades. Long-term infections increase the risk of liver cancer by approximately 50-fold. There is significant geographic variation in infection rates, but it is estimated that 300 to 350 million people worldwide have chronic HBV infection. In Southeast Asia, Africa, and China, >50% of the population is infected, and 8% to 15% become chronically infected.

Category and/or Source: New York Blood Center: Dr. Fred Prince's laboratory **Contact Time:** 10 minutes

58. Human Immunodeficiency Virus

(HIV-1) AIDS Virus: causes a long term infection that depresses the immune system. Infection with this virus resembles the common cold or flu, with symptoms appearing for 1-2 weeks and then becoming very mild and often undetectable for years after. During this first phase of infection, a person is said to be HIV positive. After some years (typically 3-10), the virus overrides the hosts immune system and kills T4 helper T cells, rendering the host susceptible to a variety of opportunistic infections. The stage of HIV infection when T cells are substantially depleted is called acquired immunodeficiency syndrome, or AIDS. Death from HIV infection is always due to infection by another, usually opportunistic, pathogen. HIV infection is very common and is increasing globally, though rates of infection in the United States have declined in the last decade. Currently, it is estimated that approximately 25 million people are infected with HIV.

Category and/or Source: UMDNJ: Dr. James Oleske's laboratory **Contact Time:** 30 seconds

59. Infectious Bovine Rhinotracheitis (IBR) Virus: causes a respiratory disease of cattle. Infection with this virus can cause secretions from the eyes, nose, and reproductive organs. It is now recognized as a cause of complex disease in cattle.

Category and/or Source: ATCC VR-188 **Contact Time:** 10 minutes

60. Influenza A/Brazil (H1N1) Virus: causes the flu in humans. Influenza viruses are known to mutate on an approximately annual basis and have potential for pandemic spread. H1N1 specifies the antigens present on the surface of the virus for that particular season/strain.

Category and/or Source: New Jersey Department of Health strain **Contact Time:** 10 minutes

61. Influenza A/Victoria (H3N2) Virus: causes the flu in humans. Influenza viruses are known to mutate on an approximately annual basis and have potential for pandemic spread. H3N2 specifies the antigens present on the surface of the virus for that particular season/strain.

Category and/or Source: ATCC VR-822, Hoffman-LaRoche, Pool # 28 **Contact Time:** 10 minutes

62. Influenza A2/Japan/305 (H2N2) Virus: causes the flu in humans. Influenza viruses are known to mutate on an approximately annual basis and have potential for pandemic spread. H2N2 specifies the antigens present on the surface of the virus for that particular season/strain.causes the flu in humans.

Category and/or Source: ATCC VR-100 **Contact Time:** 10 minutes

63. Influenza B Virus: causes the flu in humans. This virus evolves much more slowly than closely related Influenza A virus, and as such is not as significant a source of seasonal disease in humans.

Category and/or Source: Allen strain VR-102 **Contact Time:** 10 minutes

64. Influenza C Virus: causes the flu in humans. This is the most slowly evolving of the influenza viruses, and is know to infect both humans and pigs.

Category and/or Source: Taylor strain VR-104 **Contact Time:** 10 minutes

65-66. Klebsiella oxytoca: causes high fever, chills, flu-like symptoms and a cough productive of a lot of mucous in humans. This bacterium is considered opportunistic, but can be deadly once infections are established.

Category and/or Source: Gram negative clinical isolate | Antibiotic resistant gram negative rod **Contact Time:** 10 minutes


33 CLINTON ROAD • SUITE 102 • CLINTON SQUARE EXECUTIVE CENTER • WEST CALDWELL, NJ 07006 USA

VOICE: 973-575-9025 **FAX:** 973-575-9075 **E-MAIL:** INFO@MICROGENINC.COM
WEB: WWW.MICROGENINC.COM



MICROGEN

Vaccine for the Surface



67-68. *Klebsiella pneumoniae*: causes high fever, chills, flu-like symptoms and pneumonia. It can also cause gastrointestinal symptoms. This bacterium is considered opportunistic and is highly associated with hospital settings and with invasive procedures involving the airway. It is also a common cause of disease in alcoholics, presumably from aspiration of the bacteria.

Category and/or Source: Gram negative clinical isolate | Antibiotic resistant gram negative rod **Contact Time:** 10 minutes

69-70. *Klebsiella Pneumoniae type 1*: causes high fever, chills, flu-like symptoms and pneumonia. It can also cause gastrointestinal symptoms. This bacterium is considered opportunistic and is highly associated with hospital settings and with invasive procedures involving the airway. It is also a common cause of disease in alcoholics, presumably from aspiration of the bacteria.

Category and/or Source: ATCC 700603 Antibiotic resistant gram negative rod | ATCC 4352 **Contact Time:** 10 minutes

71. *Listeria monocytogenes*: causes a gastrointestinal infection in humans. This bacterium is particularly problematic as a contaminant of food. It commonly contaminates sausages and other preserved meat products. It causes diarrhea and has a tendency to infect the very old or immunocompromised.

Category and/or Source: ATCC 984 **Contact Time:** 10 minutes

72. *Malassezia pachydermatis*: causes a rare but often life-threatening fungal infection in immunocompromised humans. The fungus is common on the skin of dogs, and dogs are thought to be the major reservoir of the organism.

Category and/or Source: AMMRL (canine origin) **Contact Time:** 10 minutes

73. Measles Virus: causes a severe infection of humans that is characterized by cough, runny nose, and red eyes. A skin rash is also common. Spots inside the mouth are also indicative of this infection, but many people do not develop the spots or they are visible only briefly. Most people infected with measles recover fully, but infections are rare since vaccinations against the virus are common.

Category and/or Source: ATCC VR-24 **Contact Time:** 30 seconds

74. *Micrococcus luteus*: causes opportunistic infections in the immunocompromised in hospital settings. These bacteria are generally considered to be contaminants, but cause disease in rare instances. Notably, the bacterium is well adapted to living in or on dry environments such as the skin.

Category and/or Source: Gram positive clinical isolate **Contact Time:** 10 minutes

75-76. *Morganella morganii*: causes urinary tract infections, sepsis, pneumonia, wound infections, musculoskeletal infections, central nervous system infections, pericarditis, and spontaneous bacterial inflammation of the abdominal lining. This bacterium is a normal part of human flora and is considered to be an opportunistic pathogen.

Category and/or Source: Gram negative clinical isolate | Antibiotic resistant gram negative rod **Contact Time:** 10 minutes

77. Newcastle Disease Virus: causes a highly contagious disease in birds and occasionally causes eye infection and flu-like symptoms in highly exposed humans. This virus ranges in virulence from highly to mildly infective.

Category and/or Source: ATCC VR-109 **Contact Time:** 10 minutes

78. Parainfluenza Virus type 1: causes a disease in humans resembling a cold or the flu. This virus infects the upper airway, causing production of mucous, fever, and runny nose. In children the virus is also associated with bronchitis.

Category and/or Source: ATCC VR-105 **Contact Time:** 30 seconds

79. *Pasteurella haemolyticus*: causes a respiratory disease in cattle. Infections by this bacterium are rare.


Category and/or Source: ATCC 43823 **Contact Time:** 10 minutes

33 CLINTON ROAD • SUITE 102 • CLINTON SQUARE EXECUTIVE CENTER • WEST CALDWELL, NJ 07006 USA



MICROGEN

Vaccine for the Surface



80. Penicillium chermesinum: this fungus is thought to be a pathogen of social wasps.

Category and/or Source: Environmental fungus **Contact Time:** 10 minutes

81. Penicillium oxalicum: this fungus is a pathogen of corn that can cause allergy in humans.

Category and/or Source: Environmental fungus **Contact Time:** 5 minutes

82. Penicillium spinulosum: this fungus is a pre-harvest pathogen of sorghum.

Category and/or Source: Environmental fungus **Contact Time:** 5 minutes

83. Poliovirus type 1: causes a severe nerve infection in humans. The virus is spread by contaminated water or food by the fecal-oral route. After gastrointestinal infection, nerves are negatively affected in a percentage of cases, often resulting in paralysis. Although once common, the disease has been virtually eradicated by a successful vaccination program and by treatment of water and wastewater with disinfectants.

Category and/or Source: Chat strain **Contact Time:** 10 minutes

84. Porcine Parvovirus: causes infectious infertility in pigs. The virus infects virtually all pig herds, but infection is typically asymptomatic.

Category and/or Source: ATCC VR-742 **Contact Time:** 10 minutes

85. Porcine Respiratory & Reproductive Syndrome Virus: causes respiratory tract infection in young pigs and infertility in older pigs. The virus appeared suddenly in the Midwestern United States but has since spread worldwide.

Category and/or Source: GBL strain **Contact Time:** 10 minutes

86. Porcine Rotavirus: causes gastrointestinal infections in pigs. The virus is very similar to human rotavirus, which primarily infects children and is a major cause of diarrhea in the United States.

Category and/or Source: ATCC VR-893 **Contact Time:** 10 minutes

87. Proteus mirabilis: causes urinary tract problems in humans as well as bloodstream and wound infections. The bacterium produces large amounts of urease, which hydrolyzes to ammonia and makes the urine more alkaline. This can cause the kidney stones, which can lead to renal failure.

Category and/or Source: Gram negative clinical isolate **Contact Time:** 10 minutes

88. Proteus vulgaris: causes many different types of infection including urinary tract infections and wound infections, and is a common cause of sinus and respiratory infections. The bacterium is particularly difficult to eradicate in sinus and respiratory tissues.

Category and/or Source: Gram negative clinical isolate **Contact Time:** 10 minutes

89-92. Pseudomonas aeruginosa: causes many different types of infections, most of which are acquired in hospitals. Due to the ubiquitous nature of the bacterium in the environment, it is a common contaminant of environmental surfaces. It is also problematic because it has natural resistance to many disinfectants and can form biofilms on medical devices. Infection with Pseudomonas aeruginosa in patients with cystic fibrosis is often deadly over long periods of time.

Category and/or Source: AIDS patient isolate | Gram negative clinical isolate | Multiple (8) Antibiotic resistant gram negative rods | Multiple (8) Antibiotic resistant gram negative rods **Contact Time:** 10 minutes


33 CLINTON ROAD • SUITE 102 • CLINTON SQUARE EXECUTIVE CENTER • WEST CALDWELL, NJ 07006 USA

VOICE: 973-575-9025 **FAX:** 973-575-9075 **E-MAIL:** INFO@MICROGENINC.COM
WEB: WWW.MICROGENINC.COM



MICROGEN

Vaccine for the Surface



93. Pseudomonas fluorescens: causes infections related to blood transfusions and is a common environmental contaminant. These bacteria also have beneficial uses – they can be grown in culture to produce an antimicrobial compound called mupirocin which is effective against MRSA.

Category and/or Source: Gram negative clinical isolate **Contact Time:** 10 minutes

94. Pseudomonas pseudomallei: causes an infectious illness called melioidosis or Whitmore's disease that is most frequent in Southeast Asia and Northern Australia. Melioidosis is a lung infection that may involve a cavity of pus. The bacterium can also spread through the bloodstream to other parts of the body. Pseudomonas pseudomallei is found in soil, rice paddies and stagnant waters. Humans catch the disease by inhalation of contaminated dust or when soil contaminated by the bacteria comes in contact with abraded (scraped) skin.

Category and/or Source: Gram negative clinical isolate **Contact Time:** 10 minutes

95. Pseudomonas putida: causes spoilage of consumer products and grows robustly in a variety of environments. This bacterium is not known to be a human pathogen.

96. Pseudomonas stutzeri: causes primarily bacteremia (blood infection) in patients undergoing invasive medical procedures such as dialysis. These bacteria are considered to be opportunistic pathogens, and infection is quite rare.

Category and/or Source: Gram negative clinical isolate **Contact Time:** 10 minutes

97. Pseudorabies Virus: causes abortion, coughing, sneezing, fever, constipation, depression, seizures, and various other symptoms in piglets and mature pigs. Mortality in piglets less than one month of age is close to 100 percent. The virus is a type of porcine herpesvirus.

Category and/or Source: ATCC VR-135 **Contact Time:** 30 seconds

98. Respiratory Syncytial Virus (RSV): causes fever, runny nose, cough, and sometimes wheezing in young children. In general, symptoms of infection are flu-like. The virus is the most common cause of bronchitis in infants under 1 year old. By the time children reach 4 years of age, nearly all have been infected at least once with respiratory syncytial virus. Infections with this virus are rarely life-threatening.

Category and/or Source: Gram ATCC VR-26, Strain Long **Contact Time:** 10 minutes

99. Rhodococcus equi: causes a persistent bacterial pneumonia in young horses, and may become established as an endemic disease on some breeding farms. These bacteria are also a normal part of the bacterial flora of adult horses.

Category and/or Source: ATCC 6939 **Contact Time:** 10 minutes

100. Rotavirus: causes an acute, self-limiting gastrointestinal disease in humans that primarily affects children. Disease is characterized by watery diarrhea, nausea, vomiting and fever. Infections typically last for 3-8 days. In developed countries, the virus is rarely associated with mortality, but in the developing world rates of death can be quite high. Death from rotavirus infection in children is usually a result of dehydration from voluminous diarrhea.

Category and/or Source: Strain WA, obtained from the University of Ottawa, Canada **Contact Time:** 10 minutes

101-102: Salmonella choleraesuis @ 98% Organic Soil Load Tolerance/791 ppm Hard Water: causes severe gastrointestinal disease in humans. This genus of bacteria was recently reclassified to include two main species, S. enterica and S. typhi. As such, S. choleraesuis is now referred to as S. enterica serovar choleraesuis. This microorganism is a problematic contaminant of food products and most disease is transmitted by food. Undercooked poultry is a major source of infection with this bacterium.

Category and/or Source: ATCC 10708 | ATCC 19214 Antibiotic resistant gram negative rod **Contact Time:** 10 minutes

103. Salmonella typhi: causes typhoid fever in humans, which is a severe and often deadly infection that includes sustained fever as high as 40°C (104°F), profuse sweating, gastroenteritis, and diarrhea. In some cases, a rash of flat, rose-colored


33 CLINTON ROAD • SUITE 102 • CLINTON SQUARE EXECUTIVE CENTER • WEST CALDWELL, NJ 07006 USA

VOICE: 973-575-9025 **FAX:** 973-575-9075 **E-MAIL:** INFO@MICROGENINC.COM
WEB: WWW.MICROGENINC.COM



MICROGEN

Vaccine for the Surface



spots may also accompany infection. These bacteria are spread most commonly in developing countries through contaminated food or drinking water.

Category and/or Source: Gram ATCC 6539 **Contact Time:** 10 minutes

104. *Salomonella schottmuelleri*: causes enteric infection and fever in humans, characterized by profuse diarrhea, nausea, and vomiting. These bacteria are spread predominantly by contaminated food and water.

Category and/or Source: GBL strain **Contact Time:** 10 minutes

105. *Serratia marcescens*: causes conjunctivitis, keratitis, endophthalmitis, and tear duct infections in humans, where it is a normal part of the bacterial flora of the urinary tract and gastrointestinal system. This bacterium is easy to isolate and recognize in the laboratory because it grows as large, bright red colonies. It has been recognized as a contaminant of vaccines and may be resistant to some antibiotics, depending on the strain.

Category and/or Source: Gram negative clinical isolate **Contact Time:** 10 minutes

106. *Shigella dysenteriae*: causes severe gastrointestinal disease in humans, characterized by watery diarrhea, intestinal cramps, and fever. Infections with these bacteria usually last 5-7 days and can be spread easily to others via contamination of environmental surfaces. *Shigella dysenteriae* have a very low "infectious dose," meaning that only a few cells need be ingested to produce disease.

Category and/or Source: Gram GBL strain **Contact Time:** 10 minutes

107. *Sphingomonas paucimobilis*: causes a range of mostly hospital-related, non-life-threatening infections that typically are easily treated by antibiotic therapy. These strictly aerobic bacteria are naturally present in many land and water habitats.

Category and/or Source: Gram positive clinical isolate **Contact Time:** 10 minutes

108. *Staphylococcus aureus* @ 98% Organic Soil Load Tolerance/791 ppm Hard Water: causes infections and intoxication in humans. This bacterium can infect the skin, intestinal tract, wounds, and many other parts of the body, occasionally causing very serious meningitis, heart infections, and toxic shock. In addition to infections, humans can be harmed by *S. aureus* as a result of the toxins it produces when it is allowed to grow in food that is not refrigerated. Disease resulting from *Staphylococcus enterotoxin* intoxication is primarily gastrointestinal and involves profuse diarrhea, nausea, and vomiting with rapid onset for a brief period (usually 4-12 hours). In high doses, *Staphylococcus enterotoxin* is deadly. *S. aureus* is commonly part of the nasal flora of healthy individuals.

Category and/or Source: Gram positive clinical isolate | Toxic shock strain **Contact Time:** 10 minutes

109-112. *Staphylococcus aureus* (antibiotic resistant/toxic shock strains): cause infections and intoxications similar to antibiotic sensitive strains, but are much more problematic from a clinical perspective because the organisms either produce more/more potent toxins or resist the effects of a range of antibiotics. Methicillin-resistant *S. aureus* or MRSA is a critical pathogen, and some strains are now resistant to virtually all clinically available antibiotics. MRSA is responsible for a great deal of morbidity and mortality in the United States, especially among hospital patients.

Category and/or Source: Gram Toxic shock strain | AIDS patient isolate | ATCC 33591 METHICILLIN resistant | ATCC 6338 **Contact Time:** 10 minutes

113. *Staphylococcus auricularis*: causes infections of the skin, intestinal tract, wounds, and many other parts of the body, but infections are not generally as severe as with its close relative, *S. aureus*. These bacteria are members of the group called "coagulase-negative *Staphylococci*."

Category and/or Source: ATCC 33753 **Contact Time:** 10 minutes


114. *Staphylococcus capitis*: causes infections of the heart valves of adult humans and is commonly associated with bacteremia in neonates. Infections with this organism are often difficult to eradicate. These bacteria are members of the group called "coagulase-negative *Staphylococci*."

33 CLINTON ROAD • SUITE 102 • CLINTON SQUARE EXECUTIVE CENTER • WEST CALDWELL, NJ 07006 USA



MICROGEN

Vaccine for the Surface



Category and/or Source: Clinical isolate **Contact Time:** 10 minutes

115-116. Staphylococcus epidermidis: causes infection in people who are immunocompromised and in people who have indwelling catheters. Many strains produce a biofilm that allows them to adhere to the surfaces of medical prostheses. In addition, these bacteria are often resistant to many antibiotics. They live predominantly on the skin and as such are the most common contaminant in clinical laboratory tests. These bacteria are members of the group called “coagulase-negative Staphylococci.”

Category and/or Source: Gram Gram positive clinical isolate | Antibiotic resistant gram positive isolate **Contact Time:** 10 nutes

117. Staphylococcus hominis: causes infection in people who are immunocompromised and in people who have indwelling catheters. In addition, these bacteria are often resistant to many antibiotics. They live predominantly on the skin are generally considered to be non-pathogenic or opportunistically pathogenic. These bacteria are members of the group called “coagulase-negative Staphylococci.”

Category and/or Source: Gram ATCC 29885 **Contact Time:** 10 minutes

118. Staphylococcus saprophyticus: causes infection in people who are immunocompromised and in people who have indwelling catheters. In addition, these bacteria are often resistant to many antibiotics. They live predominantly on the skin are generally considered to be non-pathogenic or opportunistically pathogenic. These bacteria are members of the group called “coagulase-negative Staphylococci.”

Category and/or Source: Gram positive clinical isolate **Contact Time:** 10 minutes

119. Staphylococcus simulans: causes infection in people who are immunocompromised and in people who have indwelling catheters. In addition, these bacteria are often resistant to many antibiotics. They live predominantly on the skin are generally considered to be non-pathogenic or opportunistically pathogenic. These bacteria are members of the group called “coagulase-negative Staphylococci.”

Category and/or Source: ATCC 11631 **Contact Time:** 10 minutes

120. Stenotrophonas maltophilia: causes colonization of the skin and skin tissues of hospital patients and occasionally causes infections. Infections with this bacterium are usually of a result of growth of the organism to high levels in medical fluids.

Category and/or Source: Clinical isolate **Contact Time:** 10 minutes

121. Streptococcus hemolyticus: causes scarlet fever and rheumatic fever, which are both a result of the action of the body’s immune system after the infection has been cleared. This bacterium is an infrequent human pathogen. It is considered to be “Group A strep.”

Category and/or Source: Gram positive clinical isolate **Contact Time:** 10 minutes

122. Streptococcus equi var equi: causes a disease called strangles in horses, donkeys, and mules. In humans, infections are limited to a mild sore throat. This bacterium infects the respiratory tract of the animals, resulting in white discharge from the nose and further complications in about 10-20% of cases. Difficulty of breathing and inflamed lymph nodes are hallmarks of this disease.

Category and/or Source: Gram ATCC 33398 **Contact Time:** 10 minutes

123. Streptococcus equi var zooepidermicus: causes strangles in horses, but this variant may have greater transmissibility than the equi variant.

Category and/or Source: ATCC 43079 **Contact Time:** 10 minutes


33 CLINTON ROAD • SUITE 102 • CLINTON SQUARE EXECUTIVE CENTER • WEST CALDWELL, NJ 07006 USA

VOICE: 973-575-9025 **FAX:** 973-575-9075 **E-MAIL:** INFO@MICROGENINC.COM
WEB: WWW.MICROGENINC.COM



MICROGEN

Vaccine for the Surface



124. Streptococcus pneumoniae: causes a variety of infections in humans, including pneumonia, bronchitis, ear infections and more seriously, brain abscesses, meningitis, septic arthritis, and heart infections. These bacteria were the major cause of pneumonia in the early 1900's.

Category and/or Source: AIDS patient isolate **Contact Time:** 10 minutes

125. Streptococcus pneumoniae (PRSP): causes infections similar to those of antibiotic-sensitive *S. pneumoniae*, but treatment is made much more difficult by the organism's resistance to antibiotics.

Category and/or Source: ATCC 51915 **Contact Time:** 10 minutes

126-127. Streptococcus pyogenes: causes "Strep Throat" and skin infections in humans. If untreated by antibiotics, strep throat can cause Scarlet Fever, which is an autoimmune disease that can affect the heart. In addition, this bacterium can infect the skin, occasionally producing what is commonly referred to as "flesh eating disease," or necrotizing fasciitis.

Category and/or Source: ATCC 19615 | Bird M3 Clinical Isolate **Contact Time:** 10 minutes

128. Streptococcus salivarius: causes blood infections in people who have neutropenia, or depressed immune systems. This bacterium is similar in terms of size and shape to *S. pyogenes* but is much less pathogenic.

Category and/or Source: GBL strain **Contact Time:** 10 minutes

129. T1 bacteriophage: is a virus that infects bacteria. Phages are sometimes involved in the transfer of genes that encode toxins from one bacterium to the next.

Category and/or Source: ATCC 11303-B1 **Contact Time:** 10 minutes

130. T4 bacteriophage: is a virus that infects bacteria. Phages are sometimes involved in the transfer of genes that encode toxins from one bacterium to the next.

Category and/or Source: ATCC 11303-B4 **Contact Time:** 10 minutes

131. Transmissible Gastroenteritis (TGE) Virus: causes vomiting and diarrhea in pigs with a high rate of mortality. The virus initiates infection by destroying the villi (small finger-like structures) of the small intestine. After infection, pigs may shed the virus for 2-3 weeks.

Category and/or Source: ATCC VR-763 **Contact Time:** 10 minutes

132. Trichophyton mentagrophytes @ ~100 % Organic Soil Load Tolerance/395 ppm Hard Water: causes skin infections in humans. This fungus is responsible for "athletes foot," a persistent infection of the skin near the toes that can also infect the hair, skin, and nails.

Category and/or Source: ATCC 9533 **Contact Time:** 10 minutes

133. Ulocladium sp.: causes cutaneous infections in immunocompromised individuals and has also caused infections of the eyes. This fungus is a rare human pathogen.

Category and/or Source: Environmental fungus **Contact Time:** 5 minutes

134. Vaccinia Virus: causes cowpox in humans. Cowpox is a relatively mild skin infection that provides protective immunity against the much more serious (but recently eradicated) infection smallpox. The two viruses are very similar in terms of their size, shape, and genetic makeup.

Category and/or Source: Hoffmann LaRoche, Pool 57 **Contact Time:** 10 minutes


33 CLINTON ROAD • SUITE 102 • CLINTON SQUARE EXECUTIVE CENTER • WEST CALDWELL, NJ 07006 USA

VOICE: 973-575-9025 **FAX:** 973-575-9075 **E-MAIL:** INFO@MICROGENINC.COM
WEB: WWW.MICROGENINC.COM



MICROGEN

Vaccine for the Surface



135. Vesicular Stomatitis Virus: causes Influenza-like symptoms including headache, fever, pain on motion of eyes, malaise, nausea, pain in the limbs and back, as well as possible vesicular lesions in the mouth and on the lips and hands. The virus primarily infects cattle, but has a wide host range including humans, deer, and insects.

Category and/or Source: GBL strain **Contact Time:** 10 minutes

136. Yersinia enterocolitica: causes plague. Though once a major source of epidemics, outbreaks of this bacterium are now limited to transmission of the bacteria from the fleas of prairie dogs and other animals to humans. Approximately 10-100 cases of plague are recognized in the United States each year.

Category and/or Source: ATCC 23715 **Contact Time:** 10 minutes

ADDITIONAL ORGANISMS TO BE ADDED:

Clostridium difficile (vegetative cells): causes a gastrointestinal infection in humans. Disease is characterized by persistent diarrhea, and is associated with hospital stays. This bacterium exists in a vegetative (growing) form inside the human body, where it is highly sensitive to oxygen. After growth in the intestines, it can be excreted from the body in the form of environmentally-resistant endospores. Ironically, *C. difficile* infections often arise from administration of antibiotics intended to treat other infections. There is not currently a methodology accepted by the USEPA for testing endospores.

CA-MRSA: causes a number of diseases in humans, primarily painful, treatment resistant infections of the skin. CA-MRSA stands for Community-Acquired Methicillin-Resistant Staphylococcus aureus. As the name suggests, this particular strain is more likely to be found in the community at large than from healthcare environments, such as hospitals. In general, CA-MRSA is thought to be less pathogenic and less resistant to antibiotics than MRSA in general (which is often acquired in hospital settings).

CA-MRSA (USA 300 and 400): causes a number of diseases in humans, primarily painful, treatment resistant infections of the skin that may invade the bloodstream and result in high rates of mortality. The numbers "300" and "400" refer to specific DNA fingerprints (by PFGE analysis) of MRSA strains that have been isolated from community-acquired infections in the United States.

VRSA: causes a number of diseases in humans, primarily painful, treatment resistant infections of the skin that may invade the bloodstream and result in high rates of mortality. VRSA stands for Vancomycin-Resistant Staphylococcus aureus. Many MRSA are sensitive to the antibiotic Vancomycin, but this particular strain is not – rendering it extremely dangerous in clinical situations because treatment options are very limited.

INTERNATIONAL (NON-GLP) PUBLIC HEALTH CLAIMS.

1. Acinetobacter baumannii: *Acinetobacter baumannii* is a gram-negative, rod-shaped bacterium that is commonly found in water and soil. It is an opportunistic pathogen, and is particularly problematic in people who have been severely burned or who have been subjected to invasive medical procedures such as catheterization. Among the *Acinetobacter* genus, the species *baumannii* is the most relevant human pathogen. Most *A. baumannii* isolates are multi-drug resistant, which can make treatment of infections difficult. Recent outbreaks of *A. baumannii* have occurred in U.S. military field hospitals in Iraq and have highlighted the importance of control of this microorganism in the hospital environment, including on environmental surfaces. Colonization poses little threat to people who are healthy, but colonized health care workers and hospital visitors increase the risk of transmission to vulnerable or debilitated patients.

Category and/or Source: ATCC 19606 **Contact Time:** 10 minutes

2. Acinetobacter haemolyticus: *Acinetobacter haemolyticus* is a gram-negative, generally rod-shaped bacterium that is commonly found in water, soil and as a colonizer of mammals. It is an opportunistic pathogen, and is particularly problematic in people who have been severely burned or who have been subjected to invasive medical procedures such as catheterization. It can cause a variety of infections in susceptible hosts. At least one strain has been shown to produce shiga toxin 2.


Category and/or Source: ATCC 19002 **Contact Time:** 10 minutes

33 CLINTON ROAD • SUITE 102 • CLINTON SQUARE EXECUTIVE CENTER • WEST CALDWELL, NJ 07006 USA



MICROGEN

Vaccine for the Surface



3. *Aeromonas caviae*: *Aeromonas caviae* is a gram-negative, rod-shaped bacterium that is widely distributed in the aquatic environment. It is commonly thought to be an enteric pathogen, and can cause chronic diarrhea in immunocompromised adults or severe acute disease in children. It has been noted as a pathogen of concern in those with cancer. One of the most notable aspects of *Aeromonas caviae* is its association with contact lens wear, where it can cause keratitis.

Category and/or Source: ATCC 15468 **Contact Time:** 10 minutes

4. *Aeromonas hydrophila*: *Aeromonas hydrophila* is a gram-negative, rod-shaped bacterium that is present in virtually every fresh or brackish water environment. It is known to be an enteric pathogen, and can cause chronic diarrhea in immunocompromised adults or severe acute disease in children. It can also cause septicemia. It is sometimes thought of as a foodborne pathogen because of its strong association with shellfish.

Category and/or Source: ATCC 35654 **Contact Time:** 10 minutes

5. *Alcaligenes faecalis*: *Alcaligenes faecalis* is a gram negative, generally rod-shaped bacterium known to cause septicemia and meningitis in susceptible hosts. Members of the genus *Alcaligenes* can also be found in the respiratory tracts of cystic fibrosis patients. *Alcaligenes faecalis* is notable because of its (good) ability to convert toxic arsenite (a form of arsenic) to less toxic arsenate.

Category and/or Source: ATCC 35655 **Contact Time:** 10 minutes

6. *Citrobacter braakii*: *Citrobacter braakii* is gram-negative, rod shaped bacterium that is a member of the coliform group of bacteria. As a coliform, it is able to ferment lactose at elevated temperatures. From a clinical perspective, *Citrobacter braakii* is notable as an opportunistic pathogen. It has been known to cause septicemia in transplant patients and other immunocompromised persons. It can be found in urine, stool, wound infections, fish and animals.

Category and/or Source: ATCC 43162 **Contact Time:** 10 minutes

7. *Elizabethkingia meningoseptica*: *Elizabethkingia meningoseptica* was formerly known as *Flavobacterium meningosepticum* and is a gram-negative bacillus. It can be found in fresh and saltwater and is associated with meningitis and septicemia of neonates. The microorganism also has veterinary significance due to its capability to produce infections in reptiles. It is considered a very rare human pathogen.

Category and/or Source: ATCC 13253 **Contact Time:** 10 minutes

8. *Edwardsiella tarda*: *Edwardsiella tarda* is a gram-negative, rod shaped bacterium. It is generally considered an opportunistic pathogen but can also cause acute disease. The most notable diseases caused by *Edwardsiella tarda* are acute gastroenteritis and septicemia. In the environment, this organism is often found in or near water. It is also an important pathogen of fish.

Category and/or Source: ATCC 15947 **Contact Time:** 10 minutes

9. *Kocuria varians*: *Kocuria varians* is a gram-positive, round or cocci-shaped bacterium. Cells generally appear in clusters under the microscope. It can be found in the environment in beach sand, water, and on the skin of mammals. It has been associated with opportunistic infections, particularly those in hospitals where there is close contact between indwelling medical devices and the skin. One study in particular was able to pinpoint *Kocuria varians* as the cause of an infection associated with a ventriculoatrial shunt.

Category and/or Source: ATCC 51820 **Contact Time:** 10 minutes

10. *Leclercia adecarboxylata*: *Leclercia adecarboxylata* is a motile, gram-negative bacillus. It is an extremely rare human pathogen and is opportunistic in nature. When it does infect humans, it appears to be part of a multi-species infection. The environmental distribution of this organism is somewhat unknown at this time and it is rarely isolated from the environment.

Category and/or Source: ATCC 23216 **Contact Time:** 10 minutes


33 CLINTON ROAD • SUITE 102 • CLINTON SQUARE EXECUTIVE CENTER • WEST CALDWELL, NJ 07006 USA

VOICE: 973-575-9025 **FAX:** 973-575-9075 **E-MAIL:** INFO@MICROGENINC.COM
WEB: WWW.MICROGENINC.COM



MICROGEN

Vaccine for the Surface



11. Plesiomonas shigelloides: Plesiomonas shigelloides is a gram-negative bacillus that can be isolated from a multitude of environmental sources, especially freshwater as well as animals such as goats, cattle, pigs, monkeys and certain amphibians. It is not yet considered to be a definite source of human disease, but has been implicated by some researchers in gastrointestinal disease. This organism is suspected of being both toxic and invasive, owing to a suite of virulence factors it possesses.

Category and/or Source: ATCC 14029 **Contact Time:** 10 minutes

12. Providencia alcalifaciens: Providencia alcalifaciens is a gram-negative, motile member of the family Enterobacteriaceae. It is generally considered an opportunistic pathogen but can cause acute disease, particularly in the immunocompromised. The genus Providencia are also important nosocomial pathogens, causing infections associated with indwelling catheters, wounds, and severe burns. The primary residence of this organism is in the intestinal flora of healthy adults.

Category and/or Source: ATCC 9886 **Contact Time:** 10 minutes

13. Providencia stuartii: Providencia stuartii is a gram-negative, motile member of the family Enterobacteriaceae. It is generally considered an opportunistic pathogen but can cause acute disease, particularly in the immunocompromised. This particular species is especially problematic in the elderly and is highly associated with bloodstream infections, presumably resulting from colonization of wounds or indwelling medical devices. This organism can also be a problematic pathogen in the severely burned. The primary residence of this organism is in the intestinal flora of healthy adults.

Category and/or Source: ATCC 33672 **Contact Time:** 10 minutes

14. Providencia rettgeri: Providencia rettgeri is a gram-negative, motile member of the family Enterobacteriaceae. This organism is differentiated from many other species of Providencia by its apparent association with food animals such as chickens and reptiles. It has been shown to cause meningitis in crocodiles. It is generally considered an opportunistic pathogen for humans but can cause acute disease, particularly in the immunocompromised. The genus Providencia are important nosocomial pathogens, causing infections associated with indwelling catheters, wounds, and severe burns. The primary residence of this organism is in the intestinal flora of healthy adults.

Category and/or Source: ATCC 9250 **Contact Time:** 10 minutes

15. Serratia liquefaciens: Serratia liquefaciens is a gram-negative, rod shaped bacterium and a member of the family Enterobacteriaceae. Serratia liquefaciens is an opportunistic pathogen that is associated with hospital infections including septicemia. It is not considered to be as pathogenic as its close relative, Serratia marcescens. The natural home for this microorganism in the environment is the intestinal tracts of humans and other animals. It also has implications for aquaculture, where it has been shown to cause disease in juvenile rainbow trout.

Category and/or Source: ATCC 27592 **Contact Time:** 10 minutes

16. Shigella boydii: Shigella boydii is a gram negative bacterium belonging to the family Enterobacteriaceae. It is an important cause of human disease and is a frank pathogen. The infectious dose of all Shigella species is quite low, with ID50 values of approximately 200 cells having been reported in the literature. Shigella boydii is particularly relevant for surface disinfectants, since numerous cases of secondary disease transmissions through homes have been reported. Shigella species are also very common causes of laboratory acquired infections. The only environmental reservoir of Shigella is thought to be the intestinal tracts of humans.

Category and/or Source: ATCC 9207 **Contact Time:** 10 minutes


17. Shigella sonnei: Shigella sonnei is a gram negative bacterium belonging to the family Enterobacteriaceae. It is an important cause of human disease and is a frank pathogen. The infectious dose of all Shigella species is quite low, with ID50 values of approximately 200 cells having been reported in the literature. Shigella sonnei is arguably the most pathogenic of the four Shigella species and is a prolific toxin producer. This organism is particularly relevant from the standpoint of surface disinfection, because numerous cases of secondary disease transmission through homes have been reported in the scientific literature. Shigella species are very common causes of laboratory-acquired infections. The only environmental reservoir of Shigella is thought to be the intestinal tracts of humans, and children are particularly susceptible to infection by this pathogen.

33 CLINTON ROAD • SUITE 102 • CLINTON SQUARE EXECUTIVE CENTER • WEST CALDWELL, NJ 07006 USA



MICROGEN

Vaccine for the Surface



Category and/or Source: ATCC 29930 **Contact Time:** 10 minutes

18. *Aeromonas salmonicida*: *Aeromonas salmonicida* is a gram negative, short, rod-shaped bacterium belonging to the family Vibrionaceae. Though most of its close relatives are motile, *A. salmonicida* is not. It is a very important pathogen of fish, causing a disease called furunculosis (skin ulcers). Disease in humans caused by this microorganism has not been reported, but other members of the genus, such as *Aeromonas hydrophila* are known to cause disease in the immunocompromised.

Category and/or Source: ATCC 33658 **Contact Time:** 10 minutes

19. *Cellulosimicrobium cellulans*: *Cellulosimicrobium cellulans* (formerly known as *Oerskovia xanthineolytica*) is a gram-positive, branching bacillus that was originally isolated from soil. It is an opportunistic but infrequent pathogen of humans, where it has been reported to cause bacteremia. It is also an important pathogen of horses, causing premature foal birth and abortions. Like many environmental microorganisms, *Cellulosimicrobium* are intrinsically resistant to many different kinds of antibiotics. The primary reservoir of this opportunistic pathogen is thought to be soil.

Category and/or Source: ATCC 27402 **Contact Time:** 10 minutes

20. *Corynebacterium renale*: *Corynebacterium renale* is a facultatively anaerobic, gram-positive bacterium. It is an important veterinary pathogen, known to cause cystitis and pyelonephritis (severe urinary tract infection) in cattle. Fortunately, this pathogen is sensitive to many different antibiotics, including penicillins. The genus *Corynebacterium* is a common cause of urinary tract infections in humans, but infections with the species *renale* are rarely reported.

Category and/or Source: ATCC 19412 **Contact Time:** 10 minutes

21. *Delftia Acidovorans*: *Delftia acidovorans* is an aerobic, gram-negative, rod-shaped bacterium. It is generally regarded as a weak opportunistic pathogen. Most of the infections caused by this microorganism are associated with intravenous drug abusers and implanted medical devices. Notably, *D. acidovorans* is resistant to the class of antibiotics commonly used to treat systemic gram-negative infections (the aminoglycosides). The natural reservoir of this opportunistic pathogen in the environment is soil and water.

Category and/or Source: ATCC 43868 **Contact Time:** 10 minutes

22. *Listeria ivanovii*: *Listeria ivanovii* is a motile, gram-positive, rod-shaped facultatively anaerobic bacterium. It is a close relative of the human pathogen *Listeria monocytogenes*, but lacks a particular protein involved with cell-to-cell spread throughout the body, rendering it less pathogenic to humans. It is, however, occasionally reported as a suspected causative agent of human infections. In ruminant animals, this bacterium causes abortions, sepsis, and enteritis. This microorganism is commonly found in the environment, including in streams, wastewater, and soil.

Category and/or Source: ATCC 19119 **Contact Time:** 10 minutes

23. *Listeria welshmeri*: *Listeria welshmeri* is a motile, gram-positive, facultatively anaerobic rod-shaped bacterium. It is a close relative of the human pathogen *Listeria monocytogenes* but is different in that it is not generally considered to be a human pathogen. This bacterium can be found in decaying plants, sewage, soil, and water.

Category and/or Source: ATCC 35897 **Contact Time:** 10 minutes

24. *Shewanella putrefaciens*: *Shewanella putrefaciens* is a gram-negative, rod-shaped facultative anaerobe. It has been known to cause disease in humans and is generally considered to be an opportunistic pathogen. Infections with *Shewanella putrefaciens* often manifest as bacteremia or soft tissue infections. This microorganism can be found in water, fish, oily foods, and soil.

Category and/or Source: ATCC 49138 **Contact Time:** 10 minutes


25. *Shigella flexneri*: *Shigella flexneri* is a gram negative bacterium belonging to the family Enterobacteriaceae. It is an important cause of human disease and is a frank pathogen. The infectious dose of all *Shigella* species is quite low, with ID50 values of approximately 200 cells having been reported in the literature. *Shigella sonnei* accounts for about 60% of the

33 CLINTON ROAD • SUITE 102 • CLINTON SQUARE EXECUTIVE CENTER • WEST CALDWELL, NJ 07006 USA



MICROGEN

Vaccine for the Surface



Shigellosis seen in the United States, but *Shigella flexneri* (also called group B) accounts for virtually all other cases. *Shigella flexneri* also produces Reiter's syndrome (arthritis) in about 3% of cases. This organism is particularly relevant from the standpoint of surface disinfection, because numerous cases of secondary disease transmission through homes have been reported in the scientific literature. *Shigella* species are very common causes of laboratory-acquired infections. The only environmental reservoir of *Shigella* is thought to be the intestinal tracts of humans, and children are particularly susceptible to infection by this pathogen.

Category and/or Source: ATCC 12022 **Contact Time:** 10 minutes

26. *Shigella boydii* – *Shigella boydii* is a gram negative bacterium belonging to the family Enterobacteriaceae. It is an important cause of human disease and is a frank pathogen. The infectious dose of all *Shigella* species is quite low, with ID50 values of approximately 200 cells having been reported in the literature. *Shigella boydii* is particularly relevant for surface disinfectants, since numerous cases of secondary disease transmissions through homes have been reported. *Shigella* species are also very common causes of laboratory acquired infections. The only environmental reservoir of *Shigella* is thought to be the intestinal tracts of humans.

Category and/or Source: ATCC 9207 **Contact Time:** 10 minute

27. *Sphingobacterium multivorum*: *Sphingobacterium multivorum* is a gram-negative, aerobic, rodshaped bacterium. It is a very rare cause of human disease, but is occasionally reported. Disease from *Sphingobacterium multivorum* is characteristic of many diseases of weak opportunistic pathogens, wherein severely immunocompromised individuals may develop blood or soft tissue infections. *Sphingobacterium multivorum* is found in many environments including water, soil, and decaying plants.

Category and/or Source: ATCC 35656 **Contact Time:** 10 minutes

ADDITIONAL ORGANISMS FROM PUBLIC PLACES LABEL:

Echovirus II: causes severe infections in the very young. If an infant is infected within two weeks of birth, the prognosis is usually bleak. Infection by this virus may produce inflammation of the heart, brain and nervous system, and liver failure. Males are more likely than females to contract the disease, and it is spread predominantly in crowded, urban environments.

Legionella pneumophila: causes a severe respiratory infection in humans called Legionnaire's disease. This bacterium also produces a less severe type of disease called Pontiac Fever, which resembles the common cold. Infections are transmitted by aerosols, often from warm water sources such as contaminated hot tubs air conditioners, and decorative fountains.

Neisseria gonorrhoeae: causes infections of the urethra and of reproductive organs. These bacteria are transmitted almost entirely by sexual fluids. If infections persist, reproductive capabilities can become compromised, especially in women. *Neisseria* infections normally respond well to treatment with select antibiotics.

Salmonella choleraesuis (antibiotic resistant): causes severe gastrointestinal disease in humans, and treatment of the disease is made difficult by the organism's resistance to beta-lactam antibiotics (such as penicillin). This genus of bacteria was recently reclassified to include two main species, *S. enterica* and *S. typhi*. As such, *S. choleraesuis* is now referred to as *S. enterica* serovar *choleraesuis*. This microorganism is a problematic contaminant of food products and most disease is transmitted by food. Undercooked poultry is a major source of the microorganism, as it frequently resides in the digestive tracts of birds.

Treponema pallidum: causes the disease syphilis. Infections by this bacterium are sexually transmitted and occur in phases. In the first phase, small or large painless sores arise on and near the genitalia or mouth. In the second phase, skin rashes and fatigue are common. The third phase may be asymptomatic for years, but eventually neurological complications such as disorientation and loss of muscle control arise, and death usually follows. Fortunately, this bacterium is highly susceptible to antibiotics and infections are easily preventable.


33 CLINTON ROAD • SUITE 102 • CLINTON SQUARE EXECUTIVE CENTER • WEST CALDWELL, NJ 07006 USA

VOICE: 973-575-9025 **FAX:** 973-575-9075 **E-MAIL:** INFO@MICROGENINC.COM
WEB: WWW.MICROGENINC.COM



MICROGEN

Vaccine for the Surface



Efficacy Disclaimer:

All U.S. EPA accepted efficacy claims, are represented and relied on, only in the form of the current Master D-125, Public Places, CCX-151, Public Places Towelette labels, as posted under **White Papers**. Efficacy overviews of the respective list, are additionally inclusive, of a host of other Non-U.S. based data generating programs, including ongoing research and development studies, as well as a host of voluntary Non-GLP testing generation programs, as part of Microgen's continuing International Efficacy Stewardship Testing Program.

33 CLINTON ROAD • SUITE 102 • CLINTON SQUARE EXECUTIVE CENTER • WEST CALDWELL, NJ 07006 USA

VOICE: 973-575-9025 **FAX:** 973-575-9075 **E-MAIL:** INFO@MICROGENINC.COM
WEB: WWW.MICROGENINC.COM